# Visarin

Tiemonium Methylsulphate

**Composition**

**Visarin Tablet:** Each film coated tablet contains Tiemonium Methylsulphate INN 50 mg.
**Visarin Syrup:** Each 5 ml syrup contains Tiemonium Methylsulphate INN 10 mg.
**Visarin IM/IV Injection:** Each 2 ml ampoule contains Tiemonium Methylsulphate INN 5 mg.

**Description**

Tiemonium Methylsulphate is a quaternary ammonium anti-muscarinic agent with peripheral effect similar to those of atropine. Tiemonium Methylsulphate strengthens calcium bonding with phospholipids and proteins thus stabilizing the cell membrane of the GI tract.

**Indications**

Visarin is an antispasmodic drug. It reduces muscle spasms of the intestine, biliary system, bladder and uterus. Visarin is indicated for pain in gastrointestinal and biliary diseases such as gastroenteritis, diarrhoea, dysentery, biliary colic, enterocolitis, cholecystitis & colonopathies. It is also recommended in the treatment related to gynaecological and urological pain such as spasmodic dysmenorrhoea and urinary tract disease.

**Dosage & Administration**

**Tablet:** Recommended oral dose of Tiemonium Methylsulphate tablet is 2-6 tablets (100-300 mg) daily in divided dose.
**Syrup:**

- Children : 3 mg - 6 mg/kg body weight/day or 1.5 ml - 3 ml /kg body weight/day in 3 divided doses.
- 10 kg.wt. baby-2 teaspoonful 3 times daily.
- 15 kg.wt. baby-3 teaspoonful 3 times daily.
- Adult : 30 mg - 90 mg or 15 ml - 45 ml 3 times daily or as directed by physician.

**Injection:** Recommended injectable dose is one ampoule of Tiemonium Methylsulphate Injection 3 times daily, through Intravenous route slowly or Intramuscular route.

**Side Effects**

Tiemonium Methylsulphate INN may have some undesirable effects in certain individuals like risk of hypotension and tachycardia especially in case of injection.

**Precautions**

Precautions should be exercised for treatment of patients with disorders of the prostate, serious diseases of the kidney or liver, cardiac disorders, thyroid disorders, chronic bronchitis and certain gastrointestinal diseases. The risks of visual disturbances can make it dangerous to drive or use machines.

**Contraindications**

This drug must not be used in glaucoma (acute pain in the eyeball with disorders of vision), difficulties in urinating (disorders of the prostate or bladder).

**Use in Pregnancy & Lactation**

The use of this drug is not recommended during pregnancy & lactation, unless otherwise indicated by physician.

**Overdose**

In the case of overdose, seek medical attention.

**Storage**

Store in a cool (below 30°C) and dry place, protected from light.

**How Supplied**

**Visarin Tablet:** Each box contains 5 blister strips of 10 tablets.
**Visarin Syrup:** Each bottle containing 50 ml or 100 ml syrup.
**Visarin IM/IV Injection:** Each box contains 5 ampoules of 2 ml in blister pack.

Pharmasia

* Visarin Injection is manufactured by The IBN SINA Pharmaceutical Industry Ltd. for Pharmasia Ltd.

Manufactured by*
**Pharmasia Limited**
Gojariapara, Bhawal Mirzapur,
Gazipur, Bangladesh

P501804